

Suicide prevention strategies revisited: 10-year systematic review



Gil Zalsman, Keith Hawton, Danuta Wasserman, Kees van Heeringen, Ella Arensman, Marco Sarchiapone, Vladimir Carli, Cyril Höschl, Ran Barzilay, Judit Balazs, György Purebl, Jean Pierre Kahn, Pilar Alejandra Sáiz, Cendrine Bursztein Lipsicas, Julio Bobes, Doina Cozman, Ulrich Hegerl, Joseph Zohar

Summary

Background Many countries are developing suicide prevention strategies for which up-to-date, high-quality evidence is required. We present updated evidence for the effectiveness of suicide prevention interventions since 2005.

Methods We searched PubMed and the Cochrane Library using multiple terms related to suicide prevention for studies published between Jan 1, 2005, and Dec 31, 2014. We assessed seven interventions: public and physician education, media strategies, screening, restricting access to suicide means, treatments, and internet or hotline support. Data were extracted on primary outcomes of interest, namely suicidal behaviour (suicide, attempt, or ideation), and intermediate or secondary outcomes (treatment-seeking, identification of at-risk individuals, antidepressant prescription or use rates, or referrals). 18 suicide prevention experts from 13 European countries reviewed all articles and rated the strength of evidence using the Oxford criteria. Because the heterogeneity of populations and methodology did not permit formal meta-analysis, we present a narrative analysis.

Findings We identified 1797 studies, including 23 systematic reviews, 12 meta-analyses, 40 randomised controlled trials (RCTs), 67 cohort trials, and 22 ecological or population-based investigations. Evidence for restricting access to lethal means in prevention of suicide has strengthened since 2005, especially with regard to control of analgesics (overall decrease of 43% since 2005) and hot-spots for suicide by jumping (reduction of 86% since 2005, 79% to 91%). School-based awareness programmes have been shown to reduce suicide attempts (odds ratio [OR] 0.45, 95% CI 0.24–0.85; $p=0.014$) and suicidal ideation (0.5, 0.27–0.92; $p=0.025$). The anti-suicidal effects of clozapine and lithium have been substantiated, but might be less specific than previously thought. Effective pharmacological and psychological treatments of depression are important in prevention. Insufficient evidence exists to assess the possible benefits for suicide prevention of screening in primary care, in general public education and media guidelines. Other approaches that need further investigation include gatekeeper training, education of physicians, and internet and helpline support. The paucity of RCTs is a major limitation in the evaluation of preventive interventions.

Interpretation In the quest for effective suicide prevention initiatives, no single strategy clearly stands above the others. Combinations of evidence-based strategies at the individual level and the population level should be assessed with robust research designs.

Funding The Expert Platform on Mental Health, Focus on Depression, and the European College of Neuropsychopharmacology.

Introduction

Over 800 000 people worldwide die each year by suicide,¹ accounting for 1.4% of deaths worldwide. Suicide can occur at any point in the lifespan, and is the second most frequent, and in some countries the leading, cause of death among young people aged 15–24 years.¹ In addition, around 20–30 times as many suicide attempts occur.²

Suicide occurs because of a convergence of genetic,³ psychological,⁴ social, and cultural risk factors, combined with experiences of trauma and loss.⁵ Internal or external risk factors and the relations between them can be explained in models of suicide, such as stress–diathesis,⁶ gene–environment,⁷ and gene–environment and timing interactions.⁸

The complexity of this multifaceted phenomenon and low base rates, make research on suicide prevention highly challenging.⁹ However, the recognition of suicide

prevention as a public health priority¹⁰ and national prevention programmes have encouraged research, detection, treatment, and management of people at risk for suicide in many countries.^{11,12} A major review of the effectiveness of approaches to suicide prevention was done by Mann and colleagues in 2005.¹³ We did a systematic review using similar methodology to assess progress in suicide prevention research since that influential study.

Methods

Search strategy

We searched PubMed and the Cochrane library for all relevant English language studies published between Jan 1, 2005, and Dec 31, 2014. The initial search used the Medical Subject Headings identifiers for “suicide” (including the subheadings “suicide, attempted”, and

Lancet Psychiatry 2016

Published Online

June 8, 2016

[http://dx.doi.org/10.1016/S2215-0366\(16\)30030-X](http://dx.doi.org/10.1016/S2215-0366(16)30030-X)

See Online/Comment

[http://dx.doi.org/10.1016/S2215-0366\(16\)30068-2](http://dx.doi.org/10.1016/S2215-0366(16)30068-2)

Geha Mental Health Center and Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel (G Zalsman MD, R Barzilay MD); Division of Molecular Imaging and Neuropathology, Department of Psychiatry, Columbia University, New York, NY, USA (G Zalsman); Centre for Suicide Research, University of Oxford, Oxford, UK (K Hawton FMedSci); National Centre for Suicide Research and Prevention of Mental Ill-Health (NASP), Karolinska Institute, Stockholm, Sweden (D Wasserman MD, V Carli MD); Unit for Suicide Research, Ghent University, Ghent, Belgium (K van Heeringen MD); National Suicide Research Foundation, Department of Epidemiology and Public Health, University College Cork, Cork, Ireland (E Arensman PhD); Department of Medicine and Health Science, University of Molise, Via De Santis Campobasso and National Institute for Health, Migration and Poverty, Roma, Italy (M Sarchiapone MD); National Institute of Mental Health, Klecany, Czech Republic (C Höschl MD); Department of Developmental and Clinical Child Psychology, Institute of Psychology, Eotvos Lorand University, Budapest, Hungary (J Balazs MD); Institute of Behavioral Sciences, Semmelweis University Budapest, Budapest, Hungary (G Purebl MD); Université de Lorraine, Pôle de Psychiatrie et Psychologie Clinique, Centre Psychothérapique de Nancy-Laxou, Nancy-Laxou, France (J P Khan MD); Department of Psychiatry,

University of Oviedo, Centro de Investigación Biomédica en Red de Salud Mental, CIBERSAM Oviedo, Spain (P A Sáiz MD, J Bobes MD); Department of Community Mental Health, Faculty of Social Welfare and Health Sciences, University of Haifa, Haifa, Israel (C B Lipsicas PhD); Department of Clinical Psychology, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania (D Cozman MD); Department of Psychiatry and Psychotherapy, University of Leipzig, Leipzig, Germany (U Hegerl PhD); and Psychiatry Department, Sheba Health Center and Sackler School of Medicine, Tel Aviv University, Tel Aviv, Israel (J Zohar MD)

Correspondence to: Gil Zalsman, Geha Mental Health Center and Psychiatry Department, Tel Aviv University, Petach Tiqwa 49100, Israel
zalsman@post.tau.ac.il

Research in context

Evidence before this study

We reviewed evidence for the effectiveness of suicide prevention interventions published since Mann and colleagues' review in 2005. We searched PubMed and the Cochrane library systematically for all relevant English language studies published between Jan 1, 2005, and Dec 31, 2014. See Methods for details. We selected studies reporting completed suicide, attempted suicide, and suicidal ideation, as well as studies reporting intermediate outcomes including help-seeking behaviour or identification of at-risk individuals. This produced 164 publications, which were categorised for their level of evidence according to the Oxford criteria.

Added value of this study

Research published since 2005 strengthens the evidence base in several areas of suicide prevention. Restricting access to lethal means can clearly prevent suicide. There have been significant results of school-based awareness programmes in reducing suicide attempts and ideation. The anti-suicidal effects of clozapine and

lithium have been confirmed but may be less specific than previously thought. Effective pharmacological and psychological treatments of depression are important in prevention as well as education of physicians. There is insufficient evidence regarding possible benefits for suicide prevention of screening in primary care, in public education, and in media guidelines.

Implications of all the available evidence

Implementation of the evidence-supported methods described in this study via appropriate legislation, public and physician education and awareness has the potential to change public health strategies in suicide prevention plans, and significantly reduce the number of deaths due to suicide. More research using RCTs designed with standardised outcome measures and qualitative methods when applicable is needed to investigate public health approaches such as gatekeeper training, media regulation, internet-based intervention, and helplines, as well as in the two health-care approaches of physician education and screening in primary care.

“prevention and control”). “Suicide” was then combined with depression, health education, health promotion, public opinion, mass screening, family physicians, medical education, primary health care, antidepressive drugs, psychotherapy, schools, adolescents, methods, firearms, overdose, poisoning, gas poisoning, and mass media.

Data collection

RB and CBL reviewed abstracts and retrieved full-text articles that met inclusion criteria. Studies were selected if they reported primary outcomes of interest (completed or attempted suicide, or suicidal ideation), or if they included applicable, intermediate outcomes such as help-seeking behaviour or identification of at-risk individuals. The full text of these papers was retrieved and divided into subcategories of suicide prevention as reported in the medical literature: means restriction; treatment interventions including pharmacotherapy; psychotherapy; community and family-based interventions; follow-up and chain-of-care; education and awareness; media, telephone, or internet-based interventions; screening; and combined prevention interventions. Thereafter, all articles were sent to 18 suicide experts (the authors of this report) divided into eight review working groups. To ensure that key references were included in the review process, the identified references were sent to other known senior researchers in the field who were asked to suggest additional references. A face-to-face consensus meeting was held in Leiden, Netherlands, on Feb 12–14, 2015, at which all reports were categorised for their level of evidence according to the Oxford criteria.¹⁴ We excluded some manuscripts because of irrelevance or very low evidence. The results and conclusions of the review were

agreed by all authors. In total, four face-to-face meetings between the authors were organised to finalise the results and conclusions.

Role of funding source

The Expert Platform on Mental Health, Focus on Depression, and the European College of Neuropsychopharmacology supported this project (transportation and accommodation for meetings, English editing, and support for two research assistants) but had no influence on the contents of this report. The authors of this report had full access to all data in the study.

Results

Our literature search identified 1797 papers. Another 24 were obtained from other sources (figure). 224 papers were selected because they reported primary outcomes of interest or included applicable intermediate outcomes. These papers were assessed for eligibility, and 80 were excluded because of irrelevance or low evidence.

Heterogeneity in study methodology and in populations prevented a formal meta-analysis. We therefore present a narrative synthesis of the results in key domains of suicide prevention strategies.

30 studies addressed suicide prevention by means restriction (table 1). 14 (47%) of these 30 studies examined firearms, a common suicide method in countries where they are readily available. Availability in households increases risk of firearm suicides with pooled odds ratio (OR) of 3.24 (95% CI 2.41–4.40).^{18,19,28} Mann and colleagues¹³ concluded that firearm control legislation was associated with reduced suicides; however, more recent findings from the USA show mixed results, indicating that evidence was insufficient to determine the effectiveness of new laws, either alone or in

combination.^{15,20} Single studies investigating different populations in the USA, such as general versus high-risk,¹⁶ or severely mentally ill populations,¹⁷ as well as a combination of laws applied together,²¹ showed positive effects. Studies on the restriction of firearm availability in other countries also produced mixed results, with some reporting reduced firearm suicide incidence in men aged 20–40 years (Norway,²² Switzerland,²³ Israel²⁴) and in men and women of varying ages (New Zealand,²⁵ Austria²⁶), but usually without a major decrease in overall suicide rates or, alternatively, with just a modest method-substitution effect.²³ Studies in Australia failed to show an association between legislation and suicide in men¹⁹ or people aged 20–40 years,²⁷ a general decline in this method of suicide having begun before the legislation (but possibly following earlier firearm legislation).

Mann and colleagues¹³ identified changes to packaging of analgesics in the UK as beneficial in reducing deaths by suicide. Three studies in the UK have reinforced the evidence of the beneficial effects of smaller packets^{29–31} and three studies supported the effectiveness of withdrawing particularly toxic analgesics.^{45–47} The estimated reduction in number of deaths was 17 (95% CI –25 to –9) every 3 months in the post-intervention period and the overall reduction in number of deaths was about 43% (specific data not available) in the post-legislation period, after lowering the number of analgesic pills per pack.²⁹

Restrictions on the availability of pesticides contribute to reduced suicides in countries where this method of suicide is prevalent.¹³ The withdrawal of more toxic pesticides,³² restriction of access to these pesticides,³³ and measures related to decreasing absorption of toxic substances³⁴ are likely to reduce suicide in such countries. Safer storage of pesticides is another promising approach to suicide prevention in Sri Lanka³⁵ and India³⁶ but evidence is scarce. Little evidence is available on the prevention of hanging, except in psychiatric hospital inpatient units.³⁷ Potential initiatives have been proposed, such as anti-suicide shower heads in psychiatric units.³⁸

Three studies reported strong evidence that the erection of barriers at sites popular for jumping is useful, with an overall reduction in deaths by jumping of 86% (95% CI 79–91) and with little evidence of major substitution to other potential jumping sites.^{39–41}

Mann and colleagues¹³ concluded that the detoxification of domestic gas and the introduction of catalytic converters in cars were effective prevention methods. We found only one additional study on restricting the ease of purchase of charcoal, which suggested this approach might reduce suicides by carbon monoxide from charcoal burning.⁴²

Restricting prescriptions and sales of barbiturates,⁴³ and reducing concentration of caffeine tablets⁴⁴ decreased suicide incidence.

Since psychiatric disorders are a major risk factor for suicidal behaviour, their pharmacological treatment contributes substantially to the prevention of suicide.⁴⁸




Figure: PRISMA flow diagram

Reasonably strong evidence from randomised controlled trials (RCTs) shows that lithium is effective in reducing the risk of suicidal behaviour in people with mood disorders,^{49–51} which was supported by the findings of a large-scale naturalistic cohort study⁵² comparing lithium with valproate. A specific anti-suicidal effect of lithium was suggested in a population of people who had attempted suicide and who were treated with lithium, compared with people who had attempted suicide and were treated with placebo,⁵³ although the number of deaths was very small (three deaths *vs* no deaths on lithium). A large-scale naturalistic study⁵⁴ suggested that anticonvulsant mood stabilisers might also have protective effects against suicide attempts. Unadjusted rates were greater during treatment with divalproex than during treatment with lithium for suicide attempt that resulted in a visit to the emergency department (31.3 *vs* 10.8 deaths per 1000 person-years, $p < 0.001$), suicide attempt resulting in hospitalisation (10.5 *vs* 4.2 suicide attempts per 1000 person-years, $p < 0.001$), and suicide death (1.7 *vs* 0.7 per 1000 person-years; $p = 0.04$). Precise *p* values were not reported. After adjustment for age, sex, health plan, year of diagnosis, comorbid medical and psychiatric conditions, and concomitant use of other psychotropic drugs, risk of suicide death was 2.7 times higher (95% CI 1.1–6.3; $p = 0.03$) during treatment with divalproex than during treatment with lithium. Corresponding hazard ratios for non-fatal attempts were 1.7 (95% CI 1.2–2.3; $p = 0.002$) for attempts resulting in hospital admission and 1.8 (1.4–2.2; $P < 0.001$) for attempts diagnosed in the emergency department.^{54,55}

Clozapine is the only drug indicated in the USA for reduction of suicide risk in psychosis. A meta-analysis⁵⁶ of the effects of clozapine in comparison with other dopamine and serotonin-receptor antagonists

	Study type	Level of evidence
Firearm restrictions		
General population in USA	Systematic review ⁴⁵	2a
Men in USA	Ecological ⁴⁶	2a
People with serious mental illness	Systematic review ⁴⁷	2a-
General population	Systematic review ⁴⁸	2c
General population in Australia	Quasi-experimental ¹⁹	2c
General population in USA	Ecological ²⁰	2c
General population in USA	Ecological ²¹	2c
Men in Norway	Ecological ²²	2c
Adult population in Switzerland	Quasi-experimental ²³	2c
Adolescents in Israel	Quasi-experimental ²⁴	2c
General population in New Zealand	Quasi-experimental ²⁵	2c
General population in Austria	Quasi-experimental ²⁶	2c
Youth in Australia (15–44 years of age)	Ecological ²⁷	2c
Youth in USA (<20 years of age)	Case-control ²⁸	3b
Analgesic withdrawal		
General population in UK	Quasi-experimental ²⁹	2c
General population in UK	Quasi-experimental ³⁰	2c
General population in UK	Quasi-experimental ³¹	2c
Pesticide regulation		
General population in Sri Lanka	Ecological ³²	2c
General population in Taiwan	Ecological ³³	2c
Changes in pesticide content		
General population in Sri Lanka	Quasi-experimental ³⁴	3b
Pesticide storage		
General population in Sri Lanka	Quasi-experimental ³⁵	2c
General population in India	Cohort study ³⁶	2c
Restricting measures on hanging		
Psychiatric inpatients	Ecological ³⁷	2c
General population and population in prison and in psychiatric settings	Systematic review ³⁸	5
Erection of barriers at jumping hot-spots		
General population in New Zealand, UK, USA, Switzerland, and Canada	Meta-analysis ³⁹	2a
General population in Canada	Quasi-experimental ⁴⁰	2c
General population in Australia	Quasi-experimental ⁴¹	2c
Restricting access to charcoal		
General population in Hong Kong	Quasi-experimental ⁴²	2c
Restrictions on barbiturate sales		
General population in Denmark	Ecological ⁴³	2c
Restrictions on caffeine tablet sales		
General population in Sweden	Quasi-experimental ⁴⁴	2c

Oxford criteria from the Oxford Centre for Evidence-based Medicine (March, 2009).¹⁴

Table 1: Level of evidence (Oxford criteria) of suicide prevention by means restriction

(eg, olanzapine and risperidone) demonstrated anti-suicidal effect in schizophrenia.⁵⁶ However, in another meta-analysis,⁵⁷ quetiapine showed no specific effect compared with other dopamine antagonists on the occurrence of suicide and attempted suicide.

Large-scale ecological studies^{58–60} of antidepressants indicate that initiation of pharmacotherapy is not associated with an increased risk of suicide, while

continuation of pharmacotherapy for depression is associated with a reduced risk of suicide. SSRIs might increase suicidal thoughts, but not actual suicidal behaviour, in early-phase pharmacotherapy of depression in adults.⁶¹ However, emergence of suicidal ideation is low, and the risk–benefit ratio for pharmacotherapy for depression appears to favour its use.^{62,63} In depression, the administration of sertraline is associated with decreased suicidal ideation and behaviour but not with emergent suicidal thinking or behaviour.⁶⁴ People aged over 75 years, with depression treated with antidepressants might be at reduced risk of attempting suicide.⁶⁵ The results of one RCT suggest that SSRIs might exert a stronger effect than would norepinephrine-dopamine reuptake inhibitors on reduction of suicidal thoughts during the initial weeks of pharmacotherapy in high-risk patients with depression.⁶⁶ Ecological studies do not show increased sales of antidepressants to be associated with an increase in suicide rates.^{67–70} It remains unclear whether drugs for depression decrease or increase suicidal risk in patients with bipolar disorder.⁷¹ However, an evaluation of the relationship between changes in the prescription of antidepressants and changes in suicide prevalence found a clear inverse correlation in 29 European countries.⁷²

In children and adolescents with depression, evidence does not support avoidance of use of antidepressant medication because of increased risk of suicidal behaviour, although there is evidence to suggest an increased risk of suicidal ideation in this population.⁷³ Adding cognitive behavioural therapy (CBT) to fluoxetine might lead to less suicidal ideation and behaviour than treatment with fluoxetine alone.⁷³

Ketamine shows promising results as a potentially effective and rapid treatment of suicidal thoughts, independent of improvement in depression, and with minimal side-effects,⁷⁴ but effects on suicide attempts or death by suicide have not yet been shown, and effects on suicidal ideation longer than a few days have not been demonstrated. Electroconvulsive therapy was shown to rapidly reduce suicide risk in case series,^{75,76} but no controlled trials have been done.

In Mann and colleagues' review,¹³ CBT, dialectical behavioural therapy (DBT), problem-solving therapy, and intensive outpatient care with outreach were considered to be promising psychotherapies in suicide prevention; however, they had received insufficient support from RCTs or meta-analyses. Table 2 summarises the level of evidence in studies that investigated various treatment strategies.^{48–124} Eight studies reported that, compared with treatment-as-usual or minimal treatment, cognitive therapies such as CBT and manual-assisted cognitive therapy are effective in reducing suicidal ideation and behaviour in adolescents,^{86,126} adults (with mixed results),^{78,84} and patients with schizophrenia,^{80,87} as well as in patients with borderline personality disorder.^{81,85} Effects have mostly been found over medium-term follow-up.⁸⁴ DBT can reduce suicidal ideation and behaviour in adolescents⁷⁹

and women with borderline personality disorder.^{82,83} Preliminary trials have shown effectiveness of low-cost alternatives to classical DBT,^{90,91} as well as their acceptability for treatment for non-help-seeking suicidal patients.⁹² Group therapy with elements of CBT, DBT, and problem-solving therapy was effective in reducing self-harm for adolescents in one study⁷⁹ but showed a negative effect in another.¹²⁷ Specific elements for which evidence is weak or absent include skill development,⁸³ the creation of a so-called own reasons to live list,⁸⁸ and the use of modelling in skill development.¹²⁸ A multi-systemic therapy approach that addresses improving parenting skills, community, school and peer support, and engagement in pro-social activities was associated with a reduction of suicidal attempts when compared with hospitalisation in adolescents.¹²⁸ A variety of other treatments have shown effectiveness in reducing suicidality,^{89,93,95} although the level of evidence was relatively low. Psychosocial treatments have not shown clear effectiveness in reducing deaths by suicide, perhaps because of small sample sizes.⁷⁸ Community, family, and group-based interventions were not identified by Mann and colleagues.¹³ We found two meta-analyses^{96,97} that did not show community mental-health services for people with serious mental illness to be superior to standard management in preventing suicide. However, results of one study⁹⁷ showed greater acceptance of treatment, and proposed the possibility of reduced hospital admissions and suicide deaths. A meta-analysis conducted in Japan¹⁰⁰ showed a reduction in suicides among people aged over 75 years following a community-based intervention with screening and follow-up components based on pre-implementation versus post-implementation changes, but with no comparative condition.

Other studies assessing social support strategies in different populations and settings showed inconsistent effects on suicide attempts and ideation,^{98,99,101,102} but positive effects on depressive symptoms.^{101,102} One systematic review¹⁰³ of family-based interventions for patients with schizophrenia found no effect on suicide. RCTs evaluating family-based interventions in suicidal adolescents have consistently shown a clear decrease in suicidal ideation and suicide risk factors,^{104–106} and enhanced protective factors¹⁰⁵ compared with routine care. Finally, a brief family-based crisis intervention with suicidal adolescents in emergency room settings showed reduced psychiatric hospitalisations and suicide attempts at 3 months of follow-up.¹⁰⁷ Table 3 summarises the level of evidence in studies that investigated population-level prevention strategies.^{121–188}

Mann and colleagues¹³ reported mixed results on the efficacy of follow-up for people who attempt suicide. Since their report, several more studies have showed conflicting results. Contact interventions through sending regular postcards appeared to be effective in reducing repetition of suicidal behaviour in Iran, but not in high-income countries.^{112,113} Provision of information

and support through telephone or face-to-face contacts appeared to reduce suicides in low-income and middle-income countries, where mental health resources in the community are scarce, but paradoxically did not affect repetition of attempted suicide.^{108,109,111} Structured follow-up of people who attempt suicide decreased the

	Level of evidence	Study type
Lithium		
Unipolar and bipolar patients	1a	Meta-analysis ⁴⁹
Patients with major affective disorder	1a	Meta-analysis ⁵⁰
Patients with major affective disorder	1b	RCT ⁵³
General population in Denmark	2b	Cohort study ⁵¹
Lithium and valproic acid		
Patients with bipolar disorder, with a previous suicide attempt	1b	RCT ⁵⁵
Antiepileptics* and lithium		
Patients with bipolar disorder	2b	Cohort study (retrospective) ⁵⁴
Clozapine		
Patients with schizophrenia	1a	Meta-analysis ⁵⁶
Quetiapine and typical antipsychotics*		
Patients with schizophrenia	1a	Systematic review ⁵⁷
Antidepressants*		
Patients prescribed antidepressants	2c	Ecological ⁵⁸
Patients prescribed antidepressants	2c	Ecological ⁵⁹
Patients with major depressive disorder, prescribed antidepressants	2c	Cohort study (prospective) ⁶⁰
Patients with major depressive disorder, with past suicide attempt	2b	Cohort study ⁶²
Patients prescribed antidepressants	2c	Ecological ⁶⁸
General population in Sweden	2c	Ecological ⁶⁹
General population in Austria	2c	Ecological ⁷⁰
General population in Europe	2c	Ecological ⁷²
Patients prescribed antidepressants	3a	Systematic review ⁷⁷
Bipolar patients	3a	Systematic review ⁷¹
Ketamine		
Suicidal patients	1a–	Systematic review ⁷⁴
SSRIs		
Patients with major depressive disorder taking SSRIs	1a	Meta-review of systematic reviews ⁶¹
Patients with major depressive disorder	1b	RCT ⁶⁶
Elderly patients with major depressive disorder, with past suicide attempt	3b	Case-control (retrospective) ⁶⁵
SSRI (citalopram)		
Patients with schizophrenia and schizoaffective patients	1b	RCT ⁶³
SSRI (sertraline)		
Elderly patients with major depressive disorder	1b	RCT ⁶⁴
SSRI (fluoxetine)		
Adolescents with major depressive disorder	1b	RCT ⁷³
Electroconvulsive therapy		
Patients with major depressive disorder	2b	Cohort study ⁷⁵
Psychiatric inpatients	3b	Quasi-experimental ⁷⁶
Psychosocial interventions		
Patients with history of self-harm	1a	Meta-analysis ⁷⁸

(Table 2 continues on next page)

	Level of evidence	Study type
(Continued from previous page)		
Group therapy		
Adolescents with history of self-harm	1a-	Systematic review ⁷⁹
Adolescent psychiatric inpatients	3b	Quasi-experimental (retrospective open label) ⁸⁸
Prisoners with borderline personality disorder	4	Quasi-experimental ⁹³
Cognitive behavioural therapy		
Patients with schizophrenia	1b	RCT ⁸⁰
Patients with suicidal behaviour	2a	Meta-analysis ⁸⁴
Cognitive psychotherapy		
Patients with borderline personality disorder	1b	RCT ⁸¹
Patients with schizophrenia	3a	Systematic review ⁸⁷
Patients with borderline personality disorder	2b	RCT ⁸⁵
Dialectical behavioural therapy		
Patients with borderline personality disorder	1b	RCT ⁸²
Patients with borderline personality disorder	1b	RCT ⁸³
Patients with borderline personality disorder	4	Quasi-experimental ⁹⁰
Psychosocial interventions in clinical settings		
Suicidal adolescent and adult patients	2b	Systematic review ⁸⁶
Respite centre		
People in a suicidal crisis	3b	Quasi-experimental ⁸⁹
Group dialectical behavioural therapy		
Patients with borderline personality disorder	4	Quasi-experimental ⁹¹
Dialectical behavioural therapy (single meeting)		
Suicidal patients	4	Quasi-experimental ⁹²
Integrative psychotherapy		
People with suicidal ideation	4	Quasi-experimental (retrospective controlled study) ⁹⁴
Psychodynamic therapy		
Adult outpatients	4	Cohort study ⁹⁵
Intensive community-based case management		
Patients with severe mental disease	1a	Meta-analysis ⁹⁶
Community-based intervention		
Patients with severe mental disease	1a-	Meta-analysis ⁹⁷
Parental involvement in therapy		
Adolescent psychiatric patients	1b	RCT ⁹⁸
Adolescent psychiatric patients	1b	RCT ⁹⁹
Community-based screening and follow-up		
Elderly population in Japan	2a	Meta-analysis ¹⁰⁰
Culturally informed community-based treatment		
African American women attempting suicide	2b	RCT ¹⁰¹
Group intervention (hiking)		
People attempting suicide	2b	RCT ¹⁰²
Family-based intervention		
Patients with schizophrenia	1a	Meta-analysis ¹⁰³
Suicidal adolescents	1b	RCT ¹⁰⁴
Brief parent-adolescent intervention		
Parents of suicidal adolescents	1b	RCT ¹⁰⁵
Brief psycho-education to parents		
Parents of suicidal adolescents	1b	RCT ¹⁰⁶
Family crisis intervention		
Suicidal adolescents	2b	Quasi-experimental ¹⁰⁷

(Table 2 continues on next page)

number of repeated attempts^{110,114} and suicides^{108,115} in some but not all studies.^{116,117} Collaborative care with the involvement of primary health-care services in follow-up has been shown to be feasible, acceptable, and effective (in terms of ideation) compared with standard care.^{118,122,124} Positive results were identified in similar programmes targeting depressed and suicidal elderly patients.¹¹⁹⁻¹²¹ Finally, a cohort study¹⁹⁰ showed that where mental health services are available, especially with a combination of resources, suicidal behaviour was clearly reduced. On the other hand, no significant association was found between physician density and suicide rates.¹²³

Mann and colleagues¹³ concluded that few school-based programmes were evidence based or had been assessed in terms of the effectiveness in preventing suicidal behaviour. Over the past decade, evaluation studies involving school-based programmes have been of better quality. Systematic reviews,¹³²⁻¹³⁶ although including few RCTs, consistently indicate improved knowledge and attitudes towards suicide but no effect on actual suicidal behaviour. However, three large RCTs¹²⁹⁻¹³¹ emphasising mental health literacy, suicide risk awareness, and skills training in schools, showed significant effects on suicide attempts and ideation. At 12 month follow-up, there was a significant reduction in suicide attempts (OR 0.45, 95% CI 0.24-0.85; $p=0.014$) and severe suicidal ideation (0.50, 0.27-0.92; $p=0.025$), compared with the control group.¹³⁰ Prospective cohort studies^{137,139,191} assessing awareness programmes in schools showed inconsistent outcomes linked to suicidal behaviour. The level of evidence was 2c with one study showing positive,¹³⁷ one showing negative,¹³⁸ and one showing mixed outcomes.¹³⁹

Ecological studies have linked initiation of general public awareness campaigns to a significant increase in calls to helplines¹⁴⁰⁻¹⁴² but without reduction in suicides.¹⁴⁰ One study¹⁴³ showed reduced suicidal ideation and plans in a specific population of gay men, and another¹⁴⁴ showed reduction in suicides at short-term follow-up.

Education of primary care physicians targeting depression recognition and treatment was identified as one of the most effective interventions in lowering suicide rates.¹³ Since 2005, ecological studies in Sweden,¹⁴⁵ Hungary,¹⁴⁶ and Slovenia,¹⁴⁷ investigating programmes for general practitioners (GPs) have shown a significant increase in antidepressant use and decreased suicide rates.^{145,146} In a multi-component intervention,¹⁴⁶ the effectiveness of educational elements could not be separated from other elements of the intervention (free telephone consultations with local psychiatrists, a new depression clinic, and access to cheaper antidepressants). Since 2005, there have been no RCTs on this subject, but results in the former systematic review¹³ consistently showed the benefit of the GPs' educational activities.

Mann and colleagues¹³ reported that gatekeeper training was helpful in reducing the number of suicides,

provided that formalised roles and pathways to treatment were readily available.¹³ Since 2005, gatekeeper training has been studied in several populations, including military personnel,¹⁵³ public school staff,¹⁴⁸ peer helpers,¹⁴⁹ youth workers,¹⁵² clinicians,^{145,146,156} depressed persons,¹⁵⁴ and Indigenous people.¹⁵⁵ However, no RCT has shown that gatekeeper training alone affects suicide rates. Systematic reviews of studies in various populations,¹⁵⁰ as well as in Indigenous people in Australia, the USA, Canada, and New Zealand,¹⁵¹ mostly showed positive effects on knowledge, skills, and attitudes of trainees. In some of the studies from New Zealand, a decrease in suicide or suicidal behaviour was reported, but they lacked control groups.¹⁵¹

Two systematic reviews^{157,158} have shown an association between the media depiction of suicide and actual suicidal behaviour, although the methodological quality of the studies reviewed was limited. The effects seem to be bi-directional: detrimental in vulnerable populations, such as people who attempt suicide,¹⁶¹ but protective in the general population when emphasising positive coping.¹⁶⁰ Media blackouts or better reporting quality have been associated with decreased suicidal behaviour.^{159,160} Media participation in the development of guidelines assists in successful implementation, but the effectiveness of such guidelines in mitigating imitative suicides varies considerably.¹⁵⁷

Studies of telephone and internet services usually have relatively low levels of evidence. These studies have mainly focused on outcome measures such as acceptability of services by users,¹⁷⁸ identification of people at risk and referral to help services,¹⁷⁹ and compliance with referrals.¹⁸⁰ Other studies have identified specific effective characteristics in these interventions^{174,181} and service providers,¹⁸³ as well as service-use barriers.¹⁸² Some report reduction in suicidal ideation following interventions such as a brief mobile treatment intervention in Sri Lanka,¹⁷³ unguided online self-help,¹⁷⁵ and a telephone aftercare intervention.¹⁷⁶ Intervention is more efficient than wait-listing.¹⁷⁷

Mann and colleagues¹³ identified the need for assessment of the cost-effectiveness of screening in the general population versus identified at-risk populations in reduction of suicide, the predictive validity and reliability of specific screening tools, and the assessment of standard screening across different cultures. A large systematic review¹⁶² concluded that evidence was insufficient to determine the benefits of screening in primary care populations. Eight other studies assessed the screening of a total of 15 244 cases and controls.^{163,165,166,168–172} One RCT¹⁶³ showed no iatrogenic effects in youth suicide screening or in high-risk populations. A systematic review¹⁶⁷ showed that youth suicide screening programmes improved identification in adolescents at risk; however, the positive predictive value of subsequent suicidal behaviour in school settings was relatively low in some of these reports (range 6–33%).

	Level of evidence	Study type
(Continued from previous page)		
Brief intervention in emergency room and follow-up contact		
People attempting suicide	1b	RCT ¹⁰⁸
People attempting suicide	1b	RCT ¹⁰⁹
Integrative programme (outreach, problem solving, adherence, continuity)		
People attempting suicide	1b	RCT ¹¹⁰
Postcard intervention		
People with past self-poisoning	1b	RCT ¹¹¹
Adolescents and adults with repeated self-harm	1b	RCT ¹¹²
Adolescents with suicide risk	2b	RCT ¹¹³
Follow-up meeting		
Patients with self-harm	2b	Cohort study ¹¹⁴
Aftercare programme		
People attempting suicide (>15 years of age)	2b	Cohort study ¹¹⁵
Assertive outreach intervention		
People attempting suicide (>12 years of age)	2b	RCT ¹¹⁶
Chain of care intervention		
People attempting suicide	2b	Cohort study ¹¹⁷
Next-day appointment		
Adult suicidal patients	2b	RCT ¹¹⁸
Collaborative prevention in primary care		
Elderly patients with major depressive disorder	2b	RCT ¹¹⁹
Elderly patients with major depressive disorder and dysthymic patients	2b	RCT ¹²⁰
Management of depression in primary care		
Elderly patients with major depressive disorder	2b	RCT ¹²¹
Availability of safety-net of mental health services		
Suicide attempters	2c	Ecological ¹²²
Physician density		
Adolescents and young adults	2c	Ecological ¹²³
Collaborative assessment and management		
Adult suicidal patients	4	Cohort study ¹²⁴
Oxford criteria from the Oxford Centre for Evidence-based Medicine (March 2009) ¹⁴ *The general terms antidepressants, antiepileptics, or antipsychotics were used only when the study assessed heterogeneous groups and the neuroscience-based nomenclature ¹⁵ could not be used. RCT=randomised controlled trial.		

Table 2: Level of evidence (Oxford criteria) of suicide prevention strategies by treatment interventions

Screening in both school¹⁶⁸ and primary-care settings¹⁶⁹ was found to be effective and safe¹⁶³ in enhancing treatment referrals and service use in high-risk adolescents at long-term follow-up. A large RCT of suicide prevention programmes implemented in Europe¹⁸⁹ did not show significant effects of screening in reducing suicidal ideation and attempts.¹³⁰ However, screening for risk behaviours in addition to psychopathology was shown to add significant value in identifying European pupils with mental health problems.¹⁶⁴ In people aged over 75 years in Japan, the use of depression screening and psychiatrist follow-up lowered suicide prevalence by 61%.¹⁷¹

Some publications since 2005 have addressed the effect of combination and multi-level prevention programmes on suicide and suicide attempts (appendix).

See Online for appendix

	Level of evidence	Level of evidence
School-based programmes		
Adolescents	1b	RCT ¹²⁹
Adolescents	1b	RCT ¹³⁰
Children	1b	RCT ¹³¹
Young adults (students)	2a	Meta-analysis ¹³²
Adolescent	2a	Systematic review ¹³³
Children and adolescents	2a	Systematic review ¹³⁴
Adolescents	2a	Systematic review ¹³⁵
Adolescents	2a	Systematic review ¹³⁶
Adolescents	2b	Quasi-experimental ¹³⁷
Adolescents and adults (school staff)	2b	Quasi-experimental ¹³⁸
Adolescents	2b	Cohort study ¹³⁹
Public awareness campaigns		
Adult men in Austria	2c	Quasi-experimental ¹⁴⁰
General population in USA	2c	Quasi-experimental ¹⁴¹
General population in USA	2c	Ecological ¹⁴²
Gay men in Switzerland	2c	Quasi-experimental ¹⁴³
General population in Japan	2c	Ecological ¹⁴⁴
Primary care physicians education		
Primary care physicians in Sweden	2c	Quasi-experimental ¹⁴⁵
Primary care physicians in Hungary	2c	Quasi-experimental ¹⁴⁶
Primary care physicians in Slovenia	2c	Quasi-experimental ¹⁴⁷
Gatekeeper training		
School staff	1b	RCT ¹⁴⁸
Suicidal callers (to crisis line)	1b	RCT ¹⁴⁹
Mixed populations	2a	Systematic review ¹⁵⁰
Indigenous peoples in Australia, USA, Canada, and New Zealand	2a	Systematic review ¹⁵¹
Youth helpers in schools	2b	Quasi-experimental ¹⁵²
Counselling staff for veterans in USA	2b	Cohort study ¹⁵³
Adolescents and adults with major depressive disorder	2c	Cohort study ¹⁵⁴
Native children and adolescents in Alaska	2c	Ecological ¹⁵⁵
General population in Nuremberg	2c	Quasi-experimental ¹⁵⁶
Media reporting		
Media reporting	2a	Systematic review ¹⁵⁷
Mixed populations	2a	Systematic review ¹⁵⁸
General population in Austria	2c	Ecological ¹⁶⁰
Depressed outpatients in Taiwan	4	Quasi-experimental ¹⁶¹
Media blackout		
General population in Austria	2c	Quasi-experimental ¹⁵⁹
Screening		
Primary care patients	1a-	Systematic review ¹⁶²
Adolescents	1b	RCT ¹⁶³
Adolescents	1b	RCT ¹⁶⁴
Adolescents	1b	Quasi-experimental study ¹⁶⁵
Helpline callers	1b	Cohort study ¹⁶⁶
Adolescents	2a	Systematic review ¹⁶⁷
Adolescents	2b	Cohort study (longitudinal) ¹⁶⁸

(Table 3 continues on next page)

Discussion

The heterogeneity of strategies and outcome measures, as well as absence of good standards for evidence level in the literature, limits conclusions about the current effectiveness of suicide prevention strategies. However, there have clearly been major advances since the review by Mann and colleagues in 2005.¹³

There is now strong evidence that restricting access to lethal means is associated with a decrease in suicide and that substitution to other methods appears to be limited. This is clearly a major strategy to be integrated in national suicide prevention plans.

Data support the use of a few pharmacological interventions in suicide prevention. First, antidepressant pharmacotherapy treatment in adults is associated with reduced suicide risk, while initiation of pharmacotherapy does not lead to an exacerbation of suicide risk. In people aged over 75 years with depression, there is a clear beneficial effect of pharmacotherapy on the risk of attempted and completed suicide. In children and adolescents, increased risk of suicidal thoughts has to be taken into account when starting pharmacotherapy for depression. However, given the increased risk of suicide in untreated depression and the absence of an increased risk of suicide associated with pharmacotherapy, currently available evidence does not support the avoidance of initiation and continuation of pharmacotherapy for depression in children and adolescents. Therefore, the ongoing discussion about possible induction of suicidality in minors should not prevent physicians from prescribing SSRIs.⁷⁷ If a decision is made to use medication, then fluoxetine might be considered, given that it is recommended as first-line medication in guidelines. Second, lithium is effective in reducing the risk of suicide in people with mood disorders, possibly through decreasing aggression and impulsivity. Valproate might have similar efficacy in patients with bipolar disorder. Third, an anti-suicidal effect of clozapine in psychosis has been demonstrated; however, recent studies suggest that clozapine might not differ from risperidone or olanzapine in this respect. Studies of ketamine suggest promising rapid beneficial effects on reducing suicidal thoughts.

Case series show that electroconvulsive therapy provides a rapid relief of suicidal thoughts. Electroconvulsive therapy should therefore be considered earlier, rather than at its conventional last resort position for patients at risk.

Data support the efficacy of psychotherapies such as CBT and DBT. Psychodynamic psychotherapies have not been systematically studied. Understanding which treatment components (such as the development of a therapeutic alliance, the role of case-management and of significant others) might be effective in treatment of suicidality is an important line for future investigation.

The evidence on chain of care and follow-up is scarce and heterogeneous, leading to weak significance of the aggregate data. Follow-up of people who attempt suicide

is strongly supported by data and should be included in any national suicide prevention strategy. Findings that associate **mental health service availability** and reduced rates of suicide indicate the need for providing mental health services in national prevention initiatives.

Community and family-based interventions are not effective in preventing suicide in severely ill mental patients. Nevertheless, the ability of such interventions to promote treatment acceptance and to reduce hospitalisation and suicide should be noted. Although cross-cultural replication is needed, there is evidence that in elderly people, screening for depression combined with community follow-up is effective in reducing suicide risk. Family interventions with suicidal adolescents show a promising effect on suicide ideation.

RCTs increasingly show reduced suicide attempts and ideation following **school-based mental health and suicide awareness programmes**, with or without combined screening. While there has been an increase in the evaluation of general public awareness campaigns, a lack of RCTs remains a major limitation, indicating that no statements can be made about the effectiveness of these campaigns in reducing suicide.

Interventions including **training programmes for GPs** might be followed by increased prescription of antidepressants, and subsequent decrease in suicides, but a direct association between training and reduced rates is difficult to identify. Future research using RCTs should examine the value of GP training and the efficiency of their diagnostic capabilities.

No RCT showed that gatekeeper training alone affected suicide rates. Gatekeeper training is usually implemented along with other initiatives, making it difficult to identify the effect of this specific intervention on suicide rates. Intermediate outcome measures, such as referral rates and psychiatric treatment rates, should also be used.¹³ Future research should review elements of training, such as who is best to lead it, to whom it should be delivered, and with which specific content. Evaluation of the uptake of training in different specific contexts or populations is especially needed.

Although no controlled studies were done on the effect of **media** on suicidal behaviour over the last decade, a clear bidirectional effect can be established. Media should be used in collaboration with journalists as a channel for appropriate public education. Further investigation is needed on the effect of the internet and social media on suicidal behaviour.

Thus far, the evidence of telephone and internet intervention effectiveness is rather scarce and of low quality. Rapidly increasing utilisation of information and communication technologies in suicide prevention requires research assessing their efficacy.

There is insufficient evidence of the benefits of screening in primary care populations for reducing risk of suicide. If referral to treatment is the outcome measure and not suicidal behaviour, screening might be

	Level of evidence	Study type
(Continued from previous page)		
Adolescents	2b	Ecological ¹⁶⁹
Adult psychiatric outpatients	2c	Quasi-experimental ¹⁷⁰
Elderly population in Japan;	2c	Quasi-experimental ¹⁷¹
Adolescents	4	Cohort study ¹⁷²
Mobile phone intervention		
People attempting suicide	1b	RCT ¹⁷³
Telephone/internet-based intervention		
Crisis line callers/internet chat and forum users	2b	Cohort study ¹⁷⁴
Internet-based intervention		
Suicidal adults	2b	RCT ¹⁷⁵
Telephone intervention		
Adolescent outpatients (alcohol abusers)	2b	RCT ¹⁷⁶
Crisis line callers	2b	RCT ¹⁷⁷
Mobile phone intervention		
Suicide attempters	4	Cohort study ¹⁷⁸
Telephone intervention		
Suicide attempters (veterans)	4	Cohort study ¹⁷⁹
Adult crisis-line callers	4	Cohort study ¹⁸⁰
Crisis line callers	4	Cohort study ¹⁸¹
Crisis line callers	4	Cohort study ¹⁸²
Crisis line callers	5	Cohort study ¹⁸³
Combined interventions		
Elderly population in Japan	2b	Quasi-experimental ¹⁸⁴
Elderly population in Japan	2b	Quasi-experimental ¹⁸⁵
General population in Hungary	2c	Ecological ¹⁸⁶
General population in Germany	2c	Quasi-experimental ¹⁸⁷
General population in Germany	2c	Quasi-experimental ¹⁸⁸
Oxford criteria from the Oxford Centre for Evidence-based Medicine (March 2009). ¹⁴ RCT=randomised controlled trial.		

Table 3: Level of evidence (Oxford criteria) of suicide prevention using population-level prevention strategies

more effective, provided that the chain of care is continuous and useful. Even in high-risk populations, evidence does not justify the cost of expensive screening procedures.

Combinations of evidence-based strategies should be assessed on the individual and population levels using RCTs with sufficient power and similar methodologies. Future research on the efficacy of combined evidenced-based prevention strategies should focus on specific targeted populations (psychiatric patients, children and adolescents, older people, and ethnic minorities), as well as on cost-effectiveness and effect size. Data suggest that each specific risk group might need a tailored preventive approach. Priority should be given to reaching out to those who fail to seek medical or psychological help, with particular attention paid to older subjects.

The main limitation of this study is that the final decisions on the level of evidence rely on the investigators' judgments and therefore reproducibility of

the findings might be more difficult than in a formal meta-analysis. However, 18 investigators participated in this process, and decisions on disagreements were made based on consensus. There is also an inherent risk of bias at review (eg, incomplete retrieval of identified research, reporting bias). Finally, the literature review goes only to the end of 2014.

To conclude, sufficient evidence supports effective methods of prevention of suicidal behaviour. Potential interventions using new social media, mobile technologies, and **continuous monitoring of large datasets** seem to be the next field to explore in the coming decade. It is difficult to standardise methods for evaluation of suicide prevention initiatives. However, wherever possible, randomised trials should be the gold-standard approach. It is also important to look for unexpected untoward effects of initiatives, such as encouragement of suicidal behaviour by greater media attention, or method substitution where access to a common method of suicide is restricted. Evaluative research should be integral to national suicide prevention plans, including access to adequate funding to encourage and permit the necessary studies. Because suicide is a major cause of death and disability, the implementation of proven, evidence-based, and cost-effective strategies are the duty and responsibility of public health policy makers and health-care providers.

Contributors

GZ and JZ conceived the idea and designed the study. GZ drafted the first version. All authors participated in data analysis, review process, and preparation of the final version.

Declaration of interests

KH is a National Institute for Health Research (NIHR) Senior Investigator and used personal funding from NIHR to support his involvement in this work. The other authors declare no competing interests.

Acknowledgments

We thank Michaela Gerchak for editorial services.

References

- World Health Organization. WHO Suicide Data. 2015. http://www.who.int/mental_health/prevention/suicide/suicideprevent/en/ (accessed July 21, 2015).
- Wasserman D. Suicide: an unnecessary death. London: Martin Dunitz, 2001.
- van Heeringen K, Mann JJ. The neurobiology of suicide. *Lancet Psychiatry* 2014; **1**: 63–72.
- O'Connor RC, Nock MK. The psychology of suicidal behaviour. *Lancet Psychiatry* 2014; **1**: 73–85.
- Hawton K, van Heeringen K. Suicide. *Lancet* 2009; **373**: 1372–81.
- Mann JJ. Neurobiology of suicidal behaviour. *Nat Rev Neurosci* 2003; **4**: 819–28.
- Caspi A, Sugden K, Moffitt TE, et al. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 2003; **301**: 386–89.
- Zalsman G, Gutman A, Shbiro L, Rosenan R, Mann JJ, Weller A. Genetic vulnerability, timing of short-term stress and mood regulation: a rodent diffusion tensor imaging study. *Eur Neuropsychopharmacol* 2015; **25**: 2075–85.
- Turecki G, Brent DA. Suicide and suicidal behaviour. *Lancet* 2015; **387**: 1227–39.
- WHO. mhGAP Mental Health Gap Action Programme: scaling up care for mental, neurological, and substance use disorders. World Health Organization, 2008. http://www.who.int/mental_health/evidence/mhGAP (accessed March 5, 2012).
- Lapierre S, Erlangsen A, Waern M, et al. A systematic review of elderly suicide prevention programs. *Crisis* 2011; **32**: 88–98.
- van der Feltz-Cornelis CM, Sarchiapone M, Postuvan V, et al. Best practice elements of multilevel suicide prevention strategies: a review of systematic reviews. *Crisis* 2011; **32**: 319–33.
- Mann JJ, Apter A, Bertolote J, et al. Suicide prevention strategies: a systematic review. *JAMA* 2005; **294**: 2064–74.
- Oxford Centre for Evidence-based Medicine. Levels of evidence (March 2009). 2009. <http://www.cebm.net/oxford-centre-evidence-based-medicine-levels-evidence-march-2009/> (accessed Feb 1, 2016).
- Hahn RA, Bilukha O, Crosby A, et al. Firearms laws and the reduction of violence: a systematic review. *Am J Prev Med* 2005; **28**: 40–71.
- Rodriguez Andres A, Hempstead K. Gun control and suicide: the impact of state firearm regulations in the United States, 1995–2004. *Health Policy* 2011; **101**: 95–103.
- McGinty EE, Webster DW, Barry CL. Gun policy and serious mental illness: priorities for future research and policy. *Psychiatr Serv* 2014; **65**: 50–58.
- Anglemyer A, Horvath T, Rutherford G. The accessibility of firearms and risk for suicide and homicide victimization among household members: a systematic review and meta-analysis. *Ann Intern Med* 2014; **160**: 101–10.
- Klieve H, Barnes M, De Leo D. Controlling firearms use in Australia: has the 1996 gun law reform produced the decrease in rates of suicide with this method? *Soc Psychiatry Psychiatr Epidemiol* 2009; **44**: 285–92.
- Rosengart M, Cummings P, Nathens A, Heagerty P, Maier R, Rivara F. An evaluation of state firearm regulations and homicide and suicide death rates. *Inj Prev* 2005; **11**: 77–83.
- Fleegler EW, Lee LK, Monuteaux MC, Hemenway D, Mannix R. Firearm legislation and firearm-related fatalities in the United States. *JAMA Intern Med* 2013; **173**: 732–40.
- Gjertsen F, Leenaars A, Vollrath ME. Mixed impact of firearms restrictions on fatal firearm injuries in males: a national observational study. *Int J Environ Res Public Health* 2014; **11**: 487–506.
- Reisch T, Steffen T, Habenstein A, Tschacher W. Change in suicide rates in Switzerland before and after firearm restriction resulting from the 2003 “Army XXI” reform. *Am J Psychiatry* 2013; **170**: 977–84.
- Lubin G, Werbeloff N, Halperin D, Shmushkevitch M, Weiser M, Knobler HY. Decrease in suicide rates after a change of policy reducing access to firearms in adolescents: a naturalistic epidemiological study. *Suicide Life Threat Behav* 2010; **40**: 421–24.
- Beautrais AL, Fergusson DM, Horwood LJ. Firearms legislation and reductions in firearm-related suicide deaths in New Zealand. *Aust N Z J Psychiatry* 2006; **40**: 253–59.
- Kapusta ND, Etzersdorfer E, Krall C, Sonneck G. Firearm legislation reform in the European Union: impact on firearm availability, firearm suicide and homicide rates in Austria. *Br J Psychiatry* 2007; **191**: 253–57.
- McPhedran S, Baker J. Suicide prevention and method restriction: evaluating the impact of limiting access to lethal means among young Australians. *Arch Suicide Res* 2012; **16**: 135–46.
- Grossman DC, Mueller BA, Riedy C, et al. Gun storage practices and risk of youth suicide and unintentional firearm injuries. *JAMA* 2005; **293**: 707–14.
- Hawton K, Bergen H, Simkin S, et al. Long term effect of reduced pack sizes of paracetamol on poisoning deaths and liver transplant activity in England and Wales: interrupted time series analyses. *BMJ* 2013; **346**: f403.
- Hawton K, Bergen H, Simkin S, et al. Impact of different pack sizes of paracetamol in the United Kingdom and Ireland on intentional overdoses: a comparative study. *BMC Public Health* 2011; **11**: 460.
- Morgan OW, Griffiths C, Majeed A. Interrupted time-series analysis of regulations to reduce paracetamol (acetaminophen) poisoning. *PLoS Med* 2007; **4**: e105.
- Gunnell D, Fernando R, Hewagama M, Priyangika WD, Konradsen F, Eddleston M. The impact of pesticide regulations on suicide in Sri Lanka. *Int J Epidemiol* 2007; **36**: 1235–42.
- Lin JJ, Lu TH. Trends in solids/liquids poisoning suicide rates in Taiwan: a test of the substitution hypothesis. *BMC Public Health* 2011; **11**: 712.

- 34 Wilks MF, Fernando R, Ariyananda PL, et al. Improvement in survival after paraquat ingestion following introduction of a new formulation in Sri Lanka. *PLoS Med* 2008; 5: e49.
- 35 Hawton K, Ratnayake L, Simkin S, Harriss L, Scott V. Evaluation of acceptability and use of lockable storage devices for pesticides in Sri Lanka that might assist in prevention of self-poisoning. *BMC Public Health* 2009; 9: 69.
- 36 Vijayakumar L, Jeyaseelan L, Kumar S, Mohanraj R, Devika S, Manikandan S. A central storage facility to reduce pesticide suicides—a feasibility study from India. *BMC Public Health* 2013; 13: 850.
- 37 Kapur N, Hunt IM, Windfuhr K, et al. Psychiatric in-patient care and suicide in England, 1997 to 2008: a longitudinal study. *Psychol Med* 2013; 43: 61–71.
- 38 Gunnell D, Bennewith O, Hawton K, Simkin S, Kapur N. The epidemiology and prevention of suicide by hanging: a systematic review. *Int J Epidemiol* 2005; 34: 433–42.
- 39 Pirkis J, Spittal MJ, Cox G, Robinson J, Cheung YT, Studdert D. The effectiveness of structural interventions at suicide hotspots: a meta-analysis. *Int J Epidemiol* 2013; 42: 541–48.
- 40 Perron S, Burrows S, Fournier M, Perron PA, Ouellet F. Installation of a bridge barrier as a suicide prevention strategy in Montreal, Quebec, Canada. *Am J Public Health* 2013; 103: 1235–39.
- 41 Law CK, Svetlic J, De Leo D. Restricting access to a suicide hotspot does not shift the problem to another location. An experiment of two river bridges in Brisbane, Australia. *Aust N Z J Public Health* 2014; 38: 134–38.
- 42 Yip PS, Law CK, Fu KW, Law YW, Wong PW, Xu Y. Restricting the means of suicide by charcoal burning. *Br J Psychiatry* 2010; 196: 241–42.
- 43 Nordentoft M, Qin P, Helweg-Larsen K, Juel K. Restrictions in means for suicide: an effective tool in preventing suicide: the Danish experience. *Suicide Life Threat Behav* 2007; 37: 688–97.
- 44 Thelander G, Jonsson AK, Personne M, Forsberg GS, Lundqvist KM, Ahlner J. Caffeine fatalities—do sales restrictions prevent intentional intoxications? *Clin Toxicol (Phila)* 2010; 48: 354–58.
- 45 Hawton K, Bergen H, Simkin S, et al. Effect of withdrawal of co-proxamol on prescribing and deaths from drug poisoning in England and Wales: time series analysis. *BMJ* 2009; 338: b2270.
- 46 Hawton K, Bergen H, Simkin S, Wells C, Kapur N, Gunnell D. Six-year follow-up of impact of co-proxamol withdrawal in England and Wales on prescribing and deaths: time-series study. *PLoS Med* 2012; 9: e1001213.
- 47 Hawton K, Bergen H, Waters K, Murphy E, Cooper J, Kapur N. Impact of withdrawal of the analgesic Co-proxamol on nonfatal self-poisoning in the UK. *Crisis* 2011; 32: 81–87.
- 48 Wasserman D, Rihmer Z, Rujescu D, et al. The European Psychiatric Association (EPA) guidance on suicide treatment and prevention. *Eur Psychiatry* 2012; 27: 129–41.
- 49 Cipriani A, Hawton K, Stockton S, Geddes JR. Lithium in the prevention of suicide in mood disorders: updated systematic review and meta-analysis. *BMJ* 2013; 346: f3646.
- 50 Baldessarini RJ, Pompili M, Tondo L. Suicidal risk in antidepressant drug trials. *Arch Gen Psychiatry* 2006; 63: 246–48.
- 51 Kessing LV, Sondergard L, Kvist K, Andersen PK. Suicide risk in patients treated with lithium. *Arch Gen Psychiatry* 2005; 62: 860–66.
- 52 Goodwin FK, Fireman B, Simon GE, Hunkeler EM, Lee J, Revicki D. Suicide risk in bipolar disorder during treatment with lithium and divalproex. *JAMA* 2003; 290: 1467–73.
- 53 Lauterbach E, Felber W, Muller-Oerlinghausen B, et al. Adjunctive lithium treatment in the prevention of suicidal behaviour in depressive disorders: a randomised, placebo-controlled, 1-year trial. *Acta Psychiatr Scand* 2008; 118: 469–79.
- 54 Gibbons RD, Hur K, Brown CH, Mann JJ. Relationship between antiepileptic drugs and suicide attempts in patients with bipolar disorder. *Arch Gen Psychiatry* 2009; 66: 1354–60.
- 55 Oquendo MA, Galfalvy HC, Currier D, et al. Treatment of suicide attempters with bipolar disorder: a randomized clinical trial comparing lithium and valproate in the prevention of suicidal behavior. *Am J Psychiatry* 2011; 168: 1050–56.
- 56 Asenjo Lobos C, Komossa K, Rummel-Kluge C, et al. Clozapine versus other atypical antipsychotics for schizophrenia. *Cochrane Database Syst Rev* 2010; 11: CD006633.
- 57 Suttajit S, Srisurapanont M, Xia J, Suttajit S, Maneeton B, Maneeton N. Quetiapine versus typical antipsychotic medications for schizophrenia. *Cochrane Database Syst Rev* 2013; 5: CD007815.
- 58 Sondergard L, Kvist K, Lopez AG, Andersen PK, Kessing LV. Temporal changes in suicide rates for persons treated and not treated with antidepressants in Denmark during 1995–1999. *Acta Psychiatr Scand* 2006; 114: 168–76.
- 59 Sondergard L, Kvist K, Andersen PK, Kessing LV. Do antidepressants prevent suicide? *Int Clin Psychopharmacol* 2006; 21: 211–18.
- 60 Sondergard L, Lopez AG, Andersen PK, Kessing LV. Continued antidepressant treatment and suicide in patients with depressive disorder. *Arch Suicide Res* 2007; 11: 163–75.
- 61 Cipriani A, Geddes JR, Furukawa TA, Barbui C. Metareview on short-term effectiveness and safety of antidepressants for depression: an evidence-based approach to inform clinical practice. *Can J Psychiatry* 2007; 52: 553–62.
- 62 Mulder RT, Joyce PR, Frampton CM, Luty SE. Antidepressant treatment is associated with a reduction in suicidal ideation and suicide attempts. *Acta Psychiatr Scand* 2008; 118: 116–22.
- 63 Zisook S, Kasckow JW, Lanouette NM, et al. Augmentation with citalopram for suicidal ideation in middle-aged and older outpatients with schizophrenia and schizoaffective disorder who have subthreshold depressive symptoms: a randomized controlled trial. *J Clin Psychiatry* 2010; 71: 915–22.
- 64 Nelson JC, Delucchi K, Schneider L. Suicidal thinking and behavior during treatment with sertraline in late-life depression. *Am J Geriatr Psychiatry* 2007; 15: 573–80.
- 65 Barak Y, Olmer A, Aizenberg D. Antidepressants reduce the risk of suicide among elderly depressed patients. *Neuropsychopharmacology* 2006; 31: 178–81.
- 66 Grunebaum MF, Keilp JG, Ellis SP, et al. SSRI versus bupropion effects on symptom clusters in suicidal depression: post hoc analysis of a randomized clinical trial. *J Clin Psychiatry* 2013; 74: 872–79.
- 67 Grunebaum MF, Mann JJ. Safe use of SSRIs in young adults: how strong is evidence for new suicide warning? *Curr Psychiatry* 2007; 6: nihpa81089.
- 68 Castelpetra G, Morsanutto A, Pascolo-Fabrizi E, Isacson G. Antidepressant use and suicide prevention: a prescription database study in the region Friuli Venezia Giulia, Italy. *Acta Psychiatr Scand* 2008; 118: 382–88.
- 69 Isacson G, Holmgren A, Osby U, Ahlner J. Decrease in suicide among the individuals treated with antidepressants: a controlled study of antidepressants in suicide, Sweden 1995–2005. *Acta Psychiatr Scand* 2009; 120: 37–44.
- 70 Kapusta ND, Niederkrotenthaler T, Etzersdorfer E, et al. Influence of psychotherapist density and antidepressant sales on suicide rates. *Acta Psychiatr Scand* 2009; 119: 236–42.
- 71 Pacchiarotti I, Bond DJ, Baldessarini RJ, et al. The International Society for Bipolar Disorders (ISBD) task force report on antidepressant use in bipolar disorders. *Am J Psychiatry* 2013; 170: 1249–62.
- 72 Gusmao R, Quintao S, McDavid D, et al. Antidepressant utilization and suicide in Europe: an ecological multi-national study. *PLoS One* 2013; 8: e66455.
- 73 March JS, Silva S, Petrycki S, et al. The treatment for adolescents with depression study (TADS): long-term effectiveness and safety outcomes. *Arch Gen Psychiatry* 2007; 64: 1132–43.
- 74 Reinstatler L, Youssef NA. Ketamine as a potential treatment for suicidal ideation: a systematic review of the literature. *Drugs R D* 2015; 15: 37–43.
- 75 Kellner CH, Fink M, Knapp R, et al. Relief of expressed suicidal intent by ECT: a consortium for research in ECT study. *Am J Psychiatry* 2005; 162: 977–82.
- 76 Patel M, Patel S, Hardy DW, Benzies BJ, Tare V. Should electroconvulsive therapy be an early consideration for suicidal patients? *J ECT* 2006; 22: 113–15.
- 77 Moller HJ. Is there evidence for negative effects of antidepressants on suicidality in depressive patients? A systematic review. *Eur Arch Psychiatry Clin Neurosci* 2006; 256: 476–96.
- 78 Crawford MJ, Thomas O, Khan N, Kulinskaya E. Psychosocial interventions following self-harm: systematic review of their efficacy in preventing suicide. *Br J Psychiatry* 2007; 190: 11–17.

- 79 Burns J, Dudley M, Hazell P, Patton G. Clinical management of deliberate self-harm in young people: the need for evidence-based approaches to reduce repetition. *Aust N Z J Psychiatry* 2005; **39**: 121–28.
- 80 Bateman K, Hansen L, Turkington D, Kingdon D. Cognitive behavioral therapy reduces suicidal ideation in schizophrenia: results from a randomized controlled trial. *Suicide Life Threat Behav* 2007; **37**: 284–90.
- 81 Weinberg I, Gunderson JG, Hennen J, Cutter CJ, Jr. Manual assisted cognitive treatment for deliberate self-harm in borderline personality disorder patients. *J Pers Disord* 2006; **20**: 482–92.
- 82 Linehan MM, Comtois KA, Murray AM, et al. Two-year randomized controlled trial and follow-up of dialectical behavior therapy vs therapy by experts for suicidal behaviors and borderline personality disorder. *Arch Gen Psychiatry* 2006; **63**: 757–66.
- 83 Neacsiu AD, Rizvi SL, Linehan MM. Dialectical behavior therapy skills use as a mediator and outcome of treatment for borderline personality disorder. *Behav Res Ther* 2010; **48**: 832–39.
- 84 Tarrier N, Taylor K, Gooding P. Cognitive-behavioral interventions to reduce suicide behavior: a systematic review and meta-analysis. *Behav Modif* 2008; **32**: 77–108.
- 85 Morey LC, Lowmaster SE, Hopwood CJ. A pilot study of manual-assisted cognitive therapy with a therapeutic assessment augmentation for borderline personality disorder. *Psychiatry Res* 2010; **178**: 531–35.
- 86 Robinson J, Hetrick SE, Martin C. Preventing suicide in young people: systematic review. *Aust N Z J Psychiatry* 2011; **45**: 3–26.
- 87 Marshall M, Rathbone J. Early intervention for psychosis. *Cochrane Database Syst Rev* 2011; **6**: CD004718.
- 88 Esposito-Smythers C, McClung TJ, Fairlie AM. Adolescent perceptions of a suicide prevention group on an inpatient unit. *Arch Suicide Res* 2006; **10**: 265–75.
- 89 Briggs S, Webb L, Buhagiar J, Braun G. Maytree: a respite center for the suicidal: an evaluation. *Crisis* 2007; **28**: 140–47.
- 90 Andion O, Ferrer M, Matali J, et al. Effectiveness of combined individual and group dialectical behavior therapy compared to only individual dialectical behavior therapy: a preliminary study. *Psychotherapy (Chic)* 2012; **49**: 241–50.
- 91 Gutteling BM, Montagne B, Nijs M, van den Bosch LM. Dialectical behavior therapy: is outpatient group psychotherapy an effective alternative to individual psychotherapy? Preliminary conclusions. *Compr Psychiatry* 2012; **53**: 1161–68.
- 92 Ward-Ciesielski EF. An open pilot feasibility study of a brief dialectical behavior therapy skills-based intervention for suicidal individuals. *Suicide Life Threat Behav* 2013; **43**: 324–35.
- 93 Black DW, Blum N, McCormick B, Allen J. Systems training for emotional predictability and problem solving (STEPPS) group treatment for offenders with borderline personality disorder. *J Nerv Ment Dis* 2013; **201**: 124–29.
- 94 Jobes DA, Wong SA, Conrad AK, Drozd JF, Neal-Walden T. The collaborative assessment and management of suicidality versus treatment as usual: a retrospective study with suicidal outpatients. *Suicide Life Threat Behav* 2005; **35**: 483–97.
- 95 Perry JC, Bond M, Presniak MD. Alliance, reactions to treatment, and counter-transference in the process of recovery from suicidal phenomena in long-term dynamic psychotherapy. *Psychother Res* 2013; **23**: 592–605.
- 96 Dieterich M, Irving CB, Park B, Marshall M. Intensive case management for severe mental illness. *Cochrane Database Syst Rev* 2010; **10**: CD007906.
- 97 Malone D, Newron-Howes G, Simmonds S, Marriot S, Tyrer P. Community mental health teams (CMHTs) for people with severe mental illnesses and disordered personality. *Cochrane Database Syst Rev* 2007; **3**: CD000270.
- 98 King CA, Klaus N, Kramer A, Venkataraman S, Quinlan P, Gillespie B. The youth-nominated support team-version II for suicidal adolescents: a randomized controlled intervention trial. *J Consult Clin Psychol* 2009; **77**: 880–93.
- 99 King CA, Kramer A, Preuss L, Kerr DC, Weisse L, Venkataraman S. Youth-nominated support team for suicidal adolescents (version 1): a randomized controlled trial. *J Consult Clin Psychol* 2006; **74**: 199–206.
- 100 Oyama H, Sakashita T, Ono Y, Goto M, Fujita M, Koida J. Effect of community-based intervention using depression screening on elderly suicide risk: a meta-analysis of the evidence from Japan. *Community Ment Health J* 2008; **44**: 311–20.
- 101 Zhang H, Neelarambam K, Schwenke TJ, Rhodes MN, Pittman DM, Kaslow NJ. Mediators of a culturally-sensitive intervention for suicidal African American women. *J Clin Psychol Med Settings* 2013; **20**: 401–14.
- 102 Sturm J, Ploderl M, Fartacek C, et al. Physical exercise through mountain hiking in high-risk suicide patients. A randomized crossover trial. *Acta Psychiatr Scand* 2012; **126**: 467–75.
- 103 Pharoah F, Mari J, Rathbone J, Wong W. Family intervention for schizophrenia. *Cochrane Database Syst Rev* 2010; **12**: CD000088.
- 104 Diamond GS, Wintersteen MB, Brown GK, et al. Attachment-based family therapy for adolescents with suicidal ideation: a randomized controlled trial. *J Am Acad Child Adolesc Psychiatry* 2010; **49**: 122–31.
- 105 Hooven C, Walsh E, Pike KC, Herting JR. Promoting CARE: including parents in youth suicide prevention. *Fam Community Health* 2012; **35**: 225–35.
- 106 Pineda J, Dadds MR. Family intervention for adolescents with suicidal behavior: a randomized controlled trial and mediation analysis. *J Am Acad Child Adolesc Psychiatry* 2013; **52**: 851–62.
- 107 Wharff EA, Ginnis KM, Ross AM. Family-based crisis intervention with suicidal adolescents in the emergency room: a pilot study. *Soc Work* 2012; **57**: 133–43.
- 108 Fleischmann A, Bertolote JM, Wasserman D, et al. Effectiveness of brief intervention and contact for suicide attempters: a randomized controlled trial in five countries. *Bull World Health Organ* 2008; **86**: 703–09.
- 109 Bertolote JM, Fleischmann A, De Leo D, et al. Repetition of suicide attempts: data from emergency care settings in five culturally different low-and middle-income countries participating in the WHO SUPRE-MISS Study. *Crisis* 2010; **31**: 194–201.
- 110 Hvid M, Vangborg K, Sorensen HJ, Nielsen IK, Stenborg JM, Wang AG. Preventing repetition of attempted suicide—II. The Amager project, a randomized controlled trial. *Nord J Psychiatry* 2011; **65**: 292–98.
- 111 Hassanian-Moghaddam H, Sarjami S, Kolahi AA, Carter GL. Postcards in Persia: randomised controlled trial to reduce suicidal behaviours 12 months after hospital-treated self-poisoning. *Br J Psychiatry* 2011; **198**: 309–16.
- 112 Beautrais AL, Gibb SJ, Faulkner A, Fergusson DM, Mulder RT. Postcard intervention for repeat self-harm: randomised controlled trial. *Br J Psychiatry* 2010; **197**: 55–60.
- 113 Robinson J, Yuen HP, Gook S, et al. Can receipt of a regular postcard reduce suicide-related behaviour in young help seekers? A randomized controlled trial. *Early Interv Psychiatry* 2012; **6**: 145–52.
- 114 Bilen K, Pettersson H, Owe-Larsson B, et al. Can early follow-up after deliberate self-harm reduce repetition? A prospective study of 325 patients. *J Affect Disord* 2014; **152–154**: 320–25.
- 115 Pan YJ, Chang WH, Lee MB, Chen CH, Liao SC, Caine ED. Effectiveness of a nationwide aftercare program for suicide attempters. *Psychol Med* 2013; **43**: 1447–54.
- 116 Morthorst B, Krogh J, Erlangsen A, Alberdi F, Nordentoft M. Effect of assertive outreach after suicide attempt in the AID (assertive intervention for deliberate self harm) trial: randomised controlled trial. *BMJ* 2012; **345**: e4972.
- 117 Johannessen HA, Dieserud G, De Leo D, Claussen B, Zahl PH. Chain of care for patients who have attempted suicide: a follow-up study from Baerum, Norway. *BMC Public Health* 2011; **11**: 81.
- 118 Comtois KA, Jobes DA, S SOC, et al. Collaborative assessment and management of suicidality (CAMS): feasibility trial for next-day appointment services. *Depress Anxiety* 2011; **28**: 963–72.
- 119 Alexopoulos GS, Reynolds CF, 3rd, Bruce ML, et al. Reducing suicidal ideation and depression in older primary care patients: 24-month outcomes of the PROSPECT study. *Am J Psychiatry* 2009; **166**: 882–90.
- 120 Unutzer J, Tang L, Oishi S, et al. Reducing suicidal ideation in depressed older primary care patients. *J Am Geriatr Soc* 2006; **54**: 1550–56.
- 121 Gallo JJ, Morales KH, Bogner HR, et al. Long term effect of depression care management on mortality in older adults: follow-up of cluster randomized clinical trial in primary care. *BMJ* 2013; **346**: f2570.
- 122 Cooper SL, Lezotte D, Jacobellis J, Diguiseppi C. Does availability of mental health resources prevent recurrent suicidal behavior? An ecological analysis. *Suicide Life Threat Behav* 2006; **36**: 409–17.

- 123 Sher L. Does the physician density affect suicide rates among adolescents and young adults? *Int J Adolesc Med Health* 2013; 25: 315–21.
- 124 Nielsen AC, Alberdi F, Rosenbaum B. Collaborative assessment and management of suicidality method shows effect. *Dan Med Bull* 2011; 58: A4300.
- 125 Zohar J, Stahl S, Moller HJ, et al. A review of the current nomenclature for psychotropic agents and an introduction to the Neuroscience-based Nomenclature. *Eur Neuropsychopharmacol* 2015; 25: 2318–25.
- 126 Taylor LM, Oldershaw A, Richards C, Davidson K, Schmidt U, Simic M. Development and pilot evaluation of a manualized cognitive-behavioural treatment package for adolescent self-harm. *Behav Cogn Psychother* 2011; 39: 619–25.
- 127 Green JM, Wood AJ, Kerfoot MJ, et al. Group therapy for adolescents with repeated self harm: randomised controlled trial with economic evaluation. *BMJ* 2011; 342: d682.
- 128 Ougrin D, Tranah T, Leigh E, Taylor L, Asarnow JR. Practitioner review: self-harm in adolescents. *J Child Psychol Psychiatry* 2012; 53: 337–50.
- 129 Asetine Jr RH, James A, Schilling EA, Glanovsky J. Evaluating the SOS suicide prevention program: a replication and extension. *BMC Public Health* 2007; 7: 161.
- 130 Wasserman D, Hoven CW, Wasserman C, et al. School-based suicide prevention programmes: the SEYLE cluster-randomised, controlled trial. *Lancet* 2015; 385: 1536–44.
- 131 Wilcox HC, Kellam SG, Brown CH, et al. The impact of two universal randomized first- and second-grade classroom interventions on young adult suicide ideation and attempts. *Drug Alcohol Depend* 2008; 95: S60–73.
- 132 Harrod CS, Goss CW, Stallones L, DiGiuseppi C. Interventions for primary prevention of suicide in university and other post-secondary educational settings. *Cochrane Database Syst Rev* 2014; 10: CD009439.
- 133 Cusimano MD, Sameem M. The effectiveness of middle and high school-based suicide prevention programmes for adolescents: a systematic review. *Inj Prev* 2011; 17: 43–49.
- 134 Katz C, Bolton SL, Katz LY, Isaak C, Tilston-Jones T, Sareen J. A systematic review of school-based suicide prevention programs. *Depress Anxiety* 2013; 30: 1030–45.
- 135 Klimes-Dougan B, Klingbeil DA, Meller SJ. The impact of universal suicide-prevention programs on the help-seeking attitudes and behaviors of youths. *Crisis* 2013; 34: 82–97.
- 136 Robinson J, Cox G, Malone A, et al. A systematic review of school-based interventions aimed at preventing, treating, and responding to suicide-related behavior in young people. *Crisis* 2013; 34: 164–82.
- 137 Ciffone J. Suicide prevention: an analysis and replication of a curriculum-based high school program. *Soc Work* 2007; 52: 41–49.
- 138 Freedenthal S. Adolescent help-seeking and the yellow ribbon suicide prevention program: an evaluation. *Suicide Life Threat Behav* 2010; 40: 628–39.
- 139 Hooven C, Herting JR, Snedker KA. Long-term outcomes for the promoting CARE suicide prevention program. *Am J Health Behav* 2010; 34: 721–36.
- 140 Till B, Sonneck G, Baldauf G, Steiner E, Niederkrotenthaler T. Reasons to love life. Effects of a suicide-awareness campaign on the utilization of a telephone emergency line in Austria. *Crisis* 2013; 34: 382–89.
- 141 Jenner E, Jenner LW, Matthews-Sterling M, Butts JK, Williams TE. Awareness effects of a youth suicide prevention media campaign in Louisiana. *Suicide Life Threat Behav* 2010; 40: 394–406.
- 142 Oliver RJ, Spilsbury JC, Osiecki SS, Denihan WM, Zureick JL, Friedman S. Brief report: preliminary results of a suicide awareness mass media campaign in Cuyahoga County, Ohio. *Suicide Life Threat Behav* 2008; 38: 245–49.
- 143 Wang J, Hausermann M, Berrut S, Weiss MG. The impact of a depression awareness campaign on mental health literacy and mental morbidity among gay men. *J Affect Disord* 2013; 150: 306–12.
- 144 Matsubayashi T, Ueda M, Sawada Y. The effect of public awareness campaigns on suicides: evidence from Nagoya, Japan. *J Affect Disord* 2014; 152–154: 526–29.
- 145 Henriksson S, Isacson G. Increased antidepressant use and fewer suicides in Jamtland county, Sweden, after a primary care educational programme on the treatment of depression. *Acta Psychiatr Scand* 2006; 114: 159–67.
- 146 Szanto K, Kalmar S, Hendin H, Rihmer Z, Mann JJ. A suicide prevention program in a region with a very high suicide rate. *Arch Gen Psychiatry* 2007; 64: 914–20.
- 147 Roskar S, Podlesek A, Zorko M, et al. Effects of training program on recognition and management of depression and suicide risk evaluation for Slovenian primary-care physicians: follow-up study. *Croat Med J* 2010; 51: 237–42.
- 148 Wyman PA, Brown CH, Inman J, et al. Randomized trial of a gatekeeper program for suicide prevention: 1-year impact on secondary school staff. *J Consult Clin Psychol* 2008; 76: 104–15.
- 149 Gould MS, Cross W, Pisani AR, Munfakh JL, Kleinman M. Impact of applied suicide intervention skills training on the national suicide prevention lifeline. *Suicide Life Threat Behav* 2013; 43: 676–91.
- 150 Isaac M, Elias B, Katz LY, et al. Gatekeeper training as a preventative intervention for suicide: a systematic review. *Can J Psychiatry* 2009; 54: 260–68.
- 151 Clifford AC, Doran CM, Tsey K. A systematic review of suicide prevention interventions targeting indigenous peoples in Australia, United States, Canada and New Zealand. *BMC Public Health* 2013; 13: 463.
- 152 Chagnon F, Houle J, Marcoux I, Renaud J. Control-group study of an intervention training program for youth suicide prevention. *Suicide Life Threat Behav* 2007; 37: 135–44.
- 153 Matthieu MM, Cross W, Batres AR, Flora CM, Knox KL. Evaluation of gatekeeper training for suicide prevention in veterans. *Arch Suicide Res* 2008; 12: 148–54.
- 154 Chamberlain PN, Goldney RD, Taylor AW, Eckert KA. Have mental health education programs influenced the mental health literacy of those with major depression and suicidal ideation? A comparison between 1998 and 2008 in South Australia. *Suicide Life Threat Behav* 2012; 42: 525–40.
- 155 May PA, Serna P, Hurt L, Debruyne LM. Outcome evaluation of a public health approach to suicide prevention in an American Indian tribal nation. *Am J Public Health* 2005; 95: 1238–44.
- 156 Hegerl U, Althaus D, Schmidtke A, Niklewski G. The alliance against depression: 2-year evaluation of a community-based intervention to reduce suicidality. *Psychol Med* 2006; 36: 1225–33.
- 157 Bohanna I, Wang X. Media guidelines for the responsible reporting of suicide: a review of effectiveness. *Crisis* 2012; 33: 190–98.
- 158 Sisask M, Varnik A. Media roles in suicide prevention: a systematic review. *Int J Environ Res Public Health* 2012; 9: 123–38.
- 159 Niederkrotenthaler T, Sonneck G. Assessing the impact of media guidelines for reporting on suicides in Austria: interrupted time series analysis. *Aust N Z J Psychiatry* 2007; 41: 419–28.
- 160 Niederkrotenthaler T, Voracek M, Herberth A, et al. Role of media reports in completed and prevented suicide: Werther v. Papageno effects. *Br J Psychiatry* 2010; 197: 234–43.
- 161 Cheng AT, Hawton K, Chen TH, et al. The influence of media reporting of a celebrity suicide on suicidal behavior in patients with a history of depressive disorder. *J Affect Disord* 2007; 103: 69–75.
- 162 O'Connor E, Gaynes BN, Burda BU, Soh C, Whitlock EP. Screening for and treatment of suicide risk relevant to primary care: a systematic review for the U.S. Preventive Services Task Force. *Ann Intern Med* 2013; 158: 741–54.
- 163 Gould MS, Marrocco FA, Kleinman M, et al. Evaluating iatrogenic risk of youth suicide screening programs: a randomized controlled trial. *JAMA* 2005; 293: 1635–43.
- 164 Kaess M, Brunner R, Parzer P, et al. Risk-behaviour screening for identifying adolescents with mental health problems in Europe. *Eur Child Adolesc Psychiatry* 2014; 23: 611–20.
- 165 King CA, Hill RM, Wynne HA, Cunningham RM. Adolescent suicide risk screening: the effect of communication about type of follow-up on adolescents' screening responses. *J Clin Child Adolesc Psychol* 2012; 41: 508–15.
- 166 Karver MS, Tarquini SJ, Caporino NE. The judgment of future suicide-related behavior: Helpline counselors' accuracy and agreement. *Crisis* 2010; 31: 272–80.

- 167 Pena JB, Caine ED. Screening as an approach for adolescent suicide prevention. *Suicide Life Threat Behav* 2006; **36**: 614–37.
- 168 Gould MS, Marrocco FA, Hoagwood K, Kleinman M, Amakawa L, Altschuler E. Service use by at-risk youths after school-based suicide screening. *J Am Acad Child Adolesc Psychiatry* 2009; **48**: 1193–201.
- 169 Gardner W, Klima J, Chisolm D, et al. Screening, triage, and referral of patients who report suicidal thought during a primary care visit. *Pediatrics* 2010; **125**: 945–52.
- 170 Lang M, Uttaro T, Caine E, Carpinello S, Felton C. Implementing routine suicide risk screening for psychiatric outpatients with serious mental disorders: II. Quantitative results. *Arch Suicide Res* 2009; **13**: 169–77.
- 171 Oyama H, Sakashita T, Hojo K, et al. A community-based survey and screening for depression in the elderly: the short-term effect on suicide risk in Japan. *Crisis* 2010; **31**: 100–08.
- 172 Haas A, Koestner B, Rosenberg J, et al. An interactive web-based method of outreach to college students at risk for suicide. *J Am Coll Health* 2008; **57**: 15–22.
- 173 Marasinghe RB, Edirippulige S, Kavanagh D, Smith A, Jiffry MT. Effect of mobile phone-based psychotherapy in suicide prevention: a randomized controlled trial in Sri Lanka. *J Telemed Telecare* 2012; **18**: 151–55.
- 174 Gilat I, Shahar G. Emotional first aid for a suicide crisis: comparison between telephonic hotline and internet. *Psychiatry* 2007; **70**: 12–18.
- 175 van Spijker BA, van Straten A, Kerkhof AJ. Effectiveness of online self-help for suicidal thoughts: results of a randomised controlled trial. *PLoS One* 2014; **9**: e90118.
- 176 Kaminer Y, Bursleson JA, Goldston DB, Burke RH. Suicidal ideation among adolescents with alcohol use disorders during treatment and aftercare. *Am J Addict* 2006; **15**: 43–49.
- 177 Rhee WK, Merbaum M, Strube MJ, Self SM. Efficacy of brief telephone psychotherapy with callers to a suicide hotline. *Suicide Life Threat Behav* 2005; **35**: 317–28.
- 178 Chen H, Mishara BL, Liu XX. A pilot study of mobile telephone message interventions with suicide attempters in China. *Crisis* 2010; **31**: 109–12.
- 179 Britton PC, Bossarte RM, Thompson C, Kemp J, Conner KR. Influences on call outcomes among veteran callers to the national veterans crisis line. *Suicide Life Threat Behav* 2013; **43**: 494–502.
- 180 Halderman BL, Eyman JR, Kerner L, Schlacks B. A paradigm for the telephonic assessment of suicidal ideation. *Suicide Life Threat Behav* 2009; **39**: 639–47.
- 181 Gould MS, Kalafat J, Harrismunfakh JL, Kleinman M. An evaluation of crisis hotline outcomes. Part 2: suicidal callers. *Suicide Life Threat Behav* 2007; **37**: 338–52.
- 182 Gould MS, Munfakh JL, Kleinman M, Lake AM. National suicide prevention lifeline: enhancing mental health care for suicidal individuals and other people in crisis. *Suicide Life Threat Behav* 2012; **42**: 22–35.
- 183 Mishara BL, Chagnon F, Daigle M, et al. Which helper behaviors and intervention styles are related to better short-term outcomes in telephone crisis intervention? Results from a silent monitoring study of calls to the U.S. 1-800-SUICIDE Network. *Suicide Life Threat Behav* 2007; **37**: 308–21.
- 184 Oyama H, Watanabe N, Ono Y, et al. Community-based suicide prevention through group activity for the elderly successfully reduced the high suicide rate for females. *Psychiatry Clin Neurosci* 2005; **59**: 337–44.
- 185 Oyama H, Ono Y, Watanabe N, et al. Local community intervention through depression screening and group activity for elderly suicide prevention. *Psychiatry Clin Neurosci* 2006; **60**: 110–14.
- 186 Szekely A, Konkoly Thege B, Mergl R, et al. How to decrease suicide rates in both genders? An effectiveness study of a community-based intervention (EAAD). *PLoS One* 2013; **8**: e75081.
- 187 Hegerl U, Mergl R, Havers I, et al. Sustainable effects on suicidality were found for the Nuremberg alliance against depression. *Eur Arch Psychiatry Clin Neurosci* 2010; **260**: 401–06.
- 188 Hubner-Liebermann B, Neuner T, Hegerl U, Hajak G, Spiessl H. Reducing suicides through an alliance against depression? *Gen Hosp Psychiatry* 2010; **32**: 514–18.
- 189 Wasserman D, Carli V, Wasserman C, et al. Saving and empowering young lives in Europe (SEYLE): a randomized controlled trial. *BMC Public Health* 2010; **10**: 192.
- 190 While D, Bickley H, Roscoe A, et al. Implementation of mental health service recommendations in England and Wales and suicide rates, 1997–2006: a cross-sectional and before-and-after observational study. *Lancet* 2012; **379**: 1005–12.
- 191 King KA, Strunk CM, Sorter MT. Preliminary effectiveness of surviving the teens suicide prevention and depression awareness program on adolescents' suicidality and self-efficacy in performing help-seeking behaviors. *J Sch Health* 2011; **81**: 581–90.